

enissov S van riel M Voit R Hekkelman M Hulsen T Hernandez N Grummt I Wehrens R and Stunnenberg H Identification of novel functional T P binding sites and general factor repertoires mbo J

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

∅

Lis JT and Kraus WL Promoter cleavage a topolIbeta and P RP collaboration ell
Ju G Lunyak V V Perissi V Garcia assets I Rose W Glass K and Rosenfeld M G topoisomerase IIbeta mediated ds N break required for regulated transcription Science

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

Fan X hou M and Struhl K ctivator specific recruitment of Mediator in vivo Nat Struct Mol iol

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

∅

∅

Iwase S Lan F ayliss P de la Torre Ubieta L Huarte M Qi H H Whetstine J R onni Roberts T M and Shi Y The X linked mental retardation gene SM X J RI defines a family of histone H lysine demethylases ell

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

Mito Y Henikoff J G and Henikoff S Histone replacement marks the boundaries of cis regulatory domains Science

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

Nelson J Santos Rosa H and Kouzarides T Proline isomerization of histone H₄ regulates lysine methylation and gene expression Cell

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

∅

Tanny J rdjument romage H Tempst P and llis Ubiquitylation of histone H₄ controls RN polymerase II transcription elongation independently of histone H₄ methylation Genes ev

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

Gomes NP jerke G Llorente Szostek S merson M and spinosa JM Gene specific requirement for P T Fb activity and RN polymerase II phosphorylation within the p transcriptional program Genes ev

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

∅

Heintzman N Stuart RK Hon G Fu Y hing W Hawkins R arrera LO Van alcar S Qu hing K Wang W Weng Z Green R rawford G and Ren istinct and predictive chromatin signatures of transcriptional promoters and enhancers in the human genome Nat Genet

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

heng S Jin VX Fan M Smith LT Liyanarachchi S Yan PS Leu YW han M W Plass Nephew KP avuluri RV and Huang TH ombinatorial analysis of transcription factor partners reveals recruitment of c MY to estrogen receptor alpha responsive promoters Mol Cell

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

Mitro N Mak P Vargas L Godio Hampton Molteni V Kreusch and Saez
The nuclear receptor LXR is a glucose sensor Nature

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

Klose R J Yan Q Tothova Z Yamane K rdjument romage H Tempst P Gilliland G Zhang
Y and Kaelin W G Jr The Retinoblastoma binding Protein R P Is an H K emethylase ell

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

ond G L and Levine J single nucleotide polymorphism in the p pathway
interacts with gender environmental stresses and tumor genetics to influence cancer in
humans Oncogene

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>

Hirano Y and Ronai Z new function for p ubiquitination ell

Le am L Linares L K Paul Julien Lacroix M Hatchi Triboulet R ossis G Shmueli
Rodriguez M S oux O and Sardet F is an atypical ubiquitin ligase that modulates p
effector functions independently of degradation ell

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve;db=PubMed;dopt=itation;list=uids>